

P -ADIC PATH SET FRACTALS

WILLIAM ABRAM AND JEFFREY C. LAGARIAS

ABSTRACT. This paper considers a class of closed p -adic sets obtained by graph-directed constructions analogous to those of Mauldin and Williams over the real numbers. These sets are characterized as those p -adic integers whose points have p -adic expansions describable by paths in the graph of a finite automaton, and we call them p -adic path set fractals. The class of such sets is shown to be closed under addition and multiplication by p -integral rational numbers $r \in \mathbb{Q} \cap \mathbb{Z}_p$, and also under set-valued addition (Minkowski sum). The Hausdorff dimensions of such sets are explicitly calculable.

1. INTRODUCTION

This paper studies closed subsets of the p -adic integers $\mathbb{Z}_p$ which are constructible in a way analogous to the (real-valued) graph-directed fractal constructions of Mauldin and Williams [20], [21]. In our p -adic context these correspond to individual sets K_v in the Mauldin-Williams construction, whose set union is the full Mauldin-Williams construction object K . We term such sets *p -adic path set fractals*.

Sets of this kind can be alternately characterized as consisting of those p -adic integers whose p -adic expansions are described by sequences generated by paths in a fixed finite automaton with a marked initial state. The collection of such paths is given abstractly in terms of one-sided symbolic dynamics, as a closed subset $X_{\mathcal{G}} \subset \mathcal{A}^{\mathbb{N}}$, where $\mathcal{A}$ is a finite symbol alphabet, specified by a presentation $\mathcal{P} = (\mathcal{G}, v)$ in which $\mathcal{G}$ is a labeled directed graph with edges labelled by elements of $\mathcal{A}$, and v is a marked initial vertex of $\mathcal{G}$. We call these objects $X_{\mathcal{G}}$ *path sets*, and study them in more detail in [2]. An important feature of path sets is that they are closed sets of $\mathcal{A}^{\mathbb{N}}$ which are generally not invariant under the (one-sided) shift $\sigma : \mathcal{A}^{\mathbb{N}} \rightarrow \mathcal{A}^{\mathbb{N}}$ given by $\sigma(\alpha_0, \alpha_1, \alpha_2, \dots) = (\alpha_1, \alpha_2, \alpha_3, \dots)$.

The first object of this paper is to relate the p -adic constructions of this paper to Mauldin-Williams real number constructions. The basic idea is that Hausdorff dimension is preserved under the map taking a p -adic expansion of a p -adic integer to the base p radix expansion of a real number, and the work is to show that the p -adic construction objects here map to particular Mauldin and Williams real number constructions. The results of Mauldin and Williams for Hausdorff dimension of (real) graph-directed fractals then yield formulas for the Hausdorff dimension of p -adic path set fractals. These use data arising from any suitable presentation $(\mathcal{G}, v)$ of a finite automaton generating the underlying path set. results.

The second, and main, object of this paper is to show that the collection of p -adic path set fractals is preserved under p -adic addition and multiplication by rational numbers r that fall in $\mathbb{Z}_p$, i.e. for those r that are p -integral. Specifically we show

that given a presentation of a p -adic path set fractal $\mathcal{P}$ one can directly compute presentations as a path set fractal of its additive translates $\mathcal{P} + r$ and multiplicative translates $r\mathcal{P}$ for all $r \in \mathbb{Q} \cap \mathbb{Z}_p$. More generally, we show that set-valued addition (Minkowski sum) of p -adic path set fractals $\mathcal{P}_1, \mathcal{P}_2$ produces another p -adic path set fractal $\mathcal{P}_1 + \mathcal{P}_2$. These results represent purely p -adic phenomena, in the sense that analogous results fail to hold over the real numbers, for Mauldin-Williams graph-directed fractal constructions.

1.1. Results. In Section 2 we review the Mauldin-Williams construction over the real numbers, then give our basic definition of p -adic path set fractals, describing them as sets specified by a solution of a set-valued functional equation using p -adic contracting maps (Theorem 2.5). We show the equivalence of this geometric definition via a p -adic Mauldin-Williams construction to the symbolic dynamics characterization using path sets. (Theorem 2.8).

Our first result deduces that p -adic path set fractals have a Hausdorff dimension that is directly computable from a suitable presentation of the underlying path set $\mathcal{P}$. A path set presentation $\mathcal{P} = (\mathcal{G}, v)$ is called *right-resolving* if the underlying directed graph $\mathcal{G}$ has the property that at each vertex of the graph, all exiting edges have different labels. All path sets have right-resolving presentations, see [2].

Theorem 1.1. (Hausdorff dimension) *Let $\mathcal{P}$ be a p -adic path set fractal, and let $(\mathcal{G}, v_0)$ be a right-resolving presentation of $\mathcal{P}$. Then its Hausdorff dimension $\alpha = d_H(\mathcal{P})$ is given by*

$$d_H(\mathcal{P}) = \log_p \sigma(A(\mathcal{G})),$$

where $\sigma(A)$ is the spectral radius of the adjacency matrix $A = A(\mathcal{G}) = [a_{i,j}]$ in which $a_{i,j}$ counts the number of directed edges from vertex i to vertex j of the underlying directed graph G of $\mathcal{G}$.

Our main results concern the effect on p -adic path set fractals of p -adic arithmetic operations. We begin with addition of p -adic rationals.

Theorem 1.2. (Closure under rational addition) *Let $\mathcal{P}$ be a p -adic path set fractal. Then for any rational number $r \in \mathbb{Q} \cap \mathbb{Z}_p$, the shifted set*

$$\mathcal{P} + r := \{x + r : x \in \mathcal{P},\}$$

is a p -adic path set fractal.

This theorem is constructive and shows that given a right-resolving presentation $\mathcal{P} = (\mathcal{G}, v)$ one can directly compute from it a right-resolving presentation $\mathcal{P} + r = (\mathcal{G}', v')$.

Theorem 1.2 also follows as a special case of the following result.

Theorem 1.3. (Closure under set addition) *Let $\mathcal{P}_1$ and $\mathcal{P}_2$ be two p -adic path set fractals in $\mathbb{Z}_p$. Then their sum-set*

$$\mathcal{P}_1 + \mathcal{P}_2 := \{x_1 + x_2 : x_1 \in \mathcal{P}_1, x_2 \in \mathcal{P}_2\}$$

is also a p -adic path set fractal.

The proof of this result shows that given right-resolving presentations of $\mathcal{P}_1 = (\mathcal{G}_1, v_1)$ and $\mathcal{P}_2 = (\mathcal{G}_2, v_2)$ one can directly construct a presentation $\mathcal{P}_1 + \mathcal{P}_2 = (\mathcal{G}_3, v_3)$, however this presentation $(\mathcal{G}_3, v_3)$ need not be right-resolving. Theorem 1.2

arises as the special case where $\mathcal{P}_2$ contains a single element, using the easy observation that the only path sets in $\mathbb{Z}_p$ consisting of a single element are those where this element is a rational $r \in \mathbb{Q} \cap \mathbb{Z}_p$ (Theorem 2.9).

Next we turn to multiplication. A main result of this paper concerns multiplication by p -adic rational numbers.

Theorem 1.4. (Closure under rational multiplication) *Let $\mathcal{P}$ be a p -adic path set fractal in $\mathbb{Q}_p$. Then for any rational number $r \in \mathbb{Q} \cap \mathbb{Z}_p$, the dilated set*

$$r\mathcal{P} := \{rx : x \in \mathcal{P}\}.$$

is also a p -adic path set fractal.

We prove this result in Section 5. This proof is constructive: given a right-resolving presentation of $\mathcal{P}$, it can produce one for $r\mathcal{P}$. Theorem 1.4 is obtained by concatenation of constructions for several special cases, as follows.

- (1) $r = M$ is a positive integer with $\gcd(p, M) = 1$. A positive integer has an infinite p -adic expansion with a finite pre-period and a periodic part with all digits 0.
- (2) $r = \frac{1}{M}$ is the inverse of a positive integer with $\gcd(p, M) = 1$;
- (3) $r = -1$. Note that -1 has a purely periodic nonterminating p -adic expansion of period 1:

$$-1 = \sum_{k=0}^{\infty} (p-1)p^k = (\cdots, p-1, p-1, p-1)_p.$$

- (4) $r = p^k$, for $k \geq 1$.

At the level of symbolic dynamics, the actions of both arithmetic operations (adding or multiplying by a p -integral rational number) applied to a p -adic path set fractal in general do *not* preserve the property of shift-invariance of the underlying path set specified by the address map for the fractal.

1.2. Extensions and generalizations. The main emphasis of this paper is on the closure of the collection of p -adic path set fractals under arithmetic operations. From the viewpoint of geometric graph-directed constructions, the closure under arithmetic operations is a special property of the p -adic numbers, and it fails to hold for Mauldin-Williams constructions made over the real numbers.

From the symbolic dynamics characterization of path sets: they are closed under set union and set intersection as well ([2, Theorem 1.2]). From these results we deduce that, given presentations of p -adic path set fractals $\mathcal{P}_1$ and $\mathcal{P}_2$ one can, in principle, compute the Hausdorff dimension of intersections of additive and multiplicative translates of these sets, such as $\mathcal{P}_1 \cap (\mathcal{P}_2 + r)$ and $\mathcal{P}_1 \cap (r\mathcal{P}_2)$. This study was motivated by questions of this kind that arose in connection with a problem of Erdős, see Erdős [9], and [15], [3]. Examples indicate that these Hausdorff dimensions can change in a complicated fashion depending on number-theoretic properties of the rational number r .

These fractals are closed under another operation: *decimation*, i.e. extracting a complete arithmetic progression of their digits. We set

$$\psi_{j,m}(\alpha_0, \alpha_1, \alpha_2, \cdots) = (\alpha_j, \alpha_{j+m}, \alpha_{j+2m}, \alpha_{j+3m}, \cdots).$$

and then define the (j, m) -decimated set

$$\psi_{j,m}(\mathcal{P}) = \{\psi_{j,m}(x) : x \in \mathcal{P}\},$$

which is itself a path set by [2, Theorem 1.5]. It might be interesting to study the effect of deimination operations on Hausdorff dimension of the images.

There are various other generalizations that might be investigated. The methods of this paper readily generalize to arithmetic operations applied to the *g-adic numbers* for arbitrary $g \geq 2$, as defined by Mahler [18]. As a topological space one has $\mathbb{Z}_g = \prod_{p|g} \mathbb{Z}_p$. However when g contains prime powers one would use a *g-adic expansion* corresponding to the alphabet $\mathcal{A} = \{0, 1, \dots, g-1\}$.

Another direction of generalization is to allow sets in the p -adic numbers $\mathbb{Q}_p$, in which case addition or multiplication of arbitrary rational numbers would be permitted. One may also generalize the notion of p -adic path sets to higher dimensions, which would correspond to $(\mathbb{Z}_p)^d$. In this case one may investigate various relaxations of the overlap conditions imposed in the Mauldin-Williams construction. In the real number analogue $\mathbb{R}^n$ results have been obtained by Ngai and Wang [23] and Das and Ngai [6].

1.3. Contents of the paper. Section 2 recalls Mauldin-Williams constructions, gives two equivalent characterizations of p -adic path set fractals, and determines such objects containing exactly one element. Section 3 gives formulas for Hausdorff dimension of path set fractals. Section 4 proves results on addition of rational numbers to p -adic path set fractals, and on set-valued addition of two p -adic path set fractals. Section 5 proves results on multiplication of p -adic path set fractals by rational numbers. Section 6 presents examples illustrating the results.

Acknowledgments. W. Abram received the support of an NSF Graduate Research Fellowship. J. Lagarias received the support of NSF grant DMS-1101373.

2. SYMBOLIC DYNAMICS AND GRAPH-DIRECTED CONSTRUCTIONS

We first review the Mauldin-Williams construction in the real number case, then formulate a p -adic analogue to it, then discuss its path set interpretation, and finally characterize those p -adic path sets containing exactly one element.

2.1. Mauldin-Williams graph-directed constructions. In the 1980's Mauldin and Williams [21] introduced general graph-directed constructions of fractal sets over the real numbers, and computed their Hausdorff dimensions, see also Edgar [7, Chap. 4]. We follow the notation established in Mauldin and Williams [21].

Definition 2.1. A *geometric graph-directed construction* in $\mathbb{R}^m$ consists of:

- (1) a finite sequence of nonoverlapping¹, compact subsets $J_1, \dots, J_n$ of $\mathbb{R}^m$, such that each J_i has nonempty interior,
- (2) a directed graph G with vertex set consisting of the integers $1, \dots, n$, such that for each pair (i, j) there is at most one directed edge from i to j . Also, for each vertex i , there is at least one exit edge, i.e. some j such that $(i, j) \in G$, and such that the underlying undirected graph is connected,
- (3) For each graph edge (i, j) there is assigned a similarity map $T_{i,j}$ of $\mathbb{R}^m$, with similarity ratio $t_{i,j}$ such that:
 - (a) for each i , $\{T_{i,j}(J_j) | (i, j) \in G\}$ is a nonoverlapping family and

$$(2.1) \quad J_i \supseteq \bigcup \{T_{i,j}(J_j) | (i, j) \in G\}$$

¹Sets J_1 and J_2 *overlap* if their intersection has nonempty interior.

- (b) if the path component of G rooted at the vertex i_1 is a cycle: $[i_1, \dots, i_q, i_{q+1} = i_1]$, then these satisfy the contraction condition

$$(2.2) \quad \prod_{k=1}^q t_{i_k, i_{k+1}} < 1.$$

Now for each i let $\mathcal{K}(J_i)$ denote the space of compact subsets of J_i , with the Hausdorff metric, ρ_H . Mauldin and Williams prove the following:

Proposition 2.2. *For each geometric graph-directed construction, there exists a unique vector of compact sets, $(K_1, \dots, K_n) \in \prod_{i=1}^n \mathcal{K}(J_i)$ such that for each i ,*

$$(2.3) \quad K_i = \bigcup \{T_{i,j}(K_j) \mid (i, j) \in G\}.$$

Proof. This is proved by Mauldin-Williams ([21, Theorem 1, p. 812]), using the results of Hutchinson [12]. Note that the maps $T_{i,j}$ act in the reverse direction to the edges of G . $\square$

The *construction object* K is then defined by

$$(2.4) \quad K = \bigcup_{i=1}^n K_i.$$

The individual K_j are the *construction sub-objects*.

Associated to the graph G is an $n \times n$ construction matrix $A = A_G$ given by

$$(2.5) \quad A = [t_{i,j}]_{1 \leq i, j \leq n},$$

where $t_{i,j}$ is defined to be zero if $(i, j) \notin G$. Now for $\beta > 0$, let

$$A_\beta = [(t_{i,j})^\beta]_{1 \leq i, j \leq n},$$

and let

$$\Phi(\beta) := \text{Spectral radius of } A_\beta.$$

Mauldin and Williams [20, Theorem 2] observe that for each construction matrix, one has

- (1) $\Phi(0) \geq 1$,
- (2) $\Phi(\beta)$ is a continuous, strictly decreasing function of $\beta \geq 0$
- (3) $\lim_{\beta \rightarrow \infty} \Phi(\beta) = 0$.

It follows that there is a unique value $\alpha \geq 1$ such that $\Phi(\alpha) = 1$, and call this the *matrix dimension*.

Mauldin and Williams determine the Hausdorff dimension of the construction object K and also of its individual sub-objects K_j , as follows. A *strongly connected component* of a directed graph G is a maximal subgraph that is strongly connected (i.e. each vertex reachable from every other vertex by a directed path.) Let $SC(G)$ denote the set of strongly connected components of the connected graph G . Put a partial ordering on $SC(G)$ by setting $H_1 \preceq H_2$ provided there is a directed path in G from a vertex in H_1 to one in H_2 . We let α_H denote the matrix dimension of the square submatrix A_H of A_G corresponding to the strongly connected component H of G , and we let C_j be the part of the graph reachable from vertex j .

Proposition 2.3. *For each geometric graph-directed construction, the Hausdorff dimension of K , the construction object, is α , where*

$$\alpha = \max\{\alpha_H \mid H \in SC(G)\}$$

and K has positive σ -finite $\mathcal{H}^\alpha$ measure.

Furthermore the Hausdorff dimension of each K_j is α_j , where

$$\alpha_j := \max\{\alpha_H | H \in C_j\},$$

and K_j has positive σ -finite α_j -dimensional Hausdorff measure. This measure is finite if and only if $\{H \in C_j | \alpha_H = \alpha_j\}$ consists of (pairwise) incomparable elements in the partial order $\preceq$ on $SC(G)$.

Proof. This is a combination of Theorems 4 and 5 of Mauldin-Williams ([21, p. 814, p. 824]). $\square$

2.2. p -adic graph-directed constructions. We formulate a (restricted) p -adic variant of the Mauldin-Williams construction inside the compact set $\mathbb{Z}_p$ as follows.

Definition 2.4. A (restricted) p -adic graph-directed construction on the p -adic integers $\mathbb{Z}_p$ consists of

- (1) a finite sequence of (identical) initial sets $J_i = \mathbb{Z}_p$, for $1 \leq i \leq n$; these sets overlap.
- (2) a finite directed labelled graph $\mathcal{G} = (G, V, E, \mathcal{L})$ with vertex set V consisting of the integers $1, 2, \dots, n$, with directed edge set E , and with labelled edge set $\mathcal{L}$, with each labeled edge assigned data $(i(e), f(e), \phi_e)$ in which $i(e), f(e) \in V$ denote the initial and final vertices of the directed edge, and the label $j_e \in \mathcal{A} = \{0, 1, \dots, p-1\}$ is drawn from the usual alphabet of p -adic digits. No two edges have the same data (i, j, ϕ_e) . Each vertex of the underlying directed graph G has at least one exit edge.
- (3) to the label j_e is associated a p -adic similarity map $\phi : \mathbb{Z}_p \rightarrow \mathbb{Z}_p$ given by

$$\phi_e(x) = px + j_e.$$

This is a contractive p -adic mapping.

We note that the conditions (a), (b) of Mauldin-Williams automatically hold, excluding the non-overlapping condition:

- (a) The initial sets $J_i = \mathbb{Z}_p$ satisfy the condition

$$J_j \supseteq \bigcup \{\phi_e(J_{i(e)}) : e \text{ has final vertex } f(e) = j\}.$$

- (b) Each map ϕ_e is has p -adic contraction ratio $t_e := |p|_p = \frac{1}{p} < 1$. Thus for $[e_1, \dots, e_q]$ a directed cycle of edges in G , the contracting cycle condition holds:

$$\prod_{i=1}^q t_{e_i} < 1,$$

This definition is broader than the Mauldin-Williams construction in that the underlying directed graph G (ignoring labels) is permitted to have loops ($i(e) = t(e)$) and multiple edges (having same $i(e), v(e)$). It differs in starting with initial sets J_i that have overlaps, which is forbidden in the Mauldin-Williams construction. On the other hand it requires that all contraction ratios be equal, and in this aspect is narrower than the Mauldin-Williams construction.

In contrast, the Mauldin-Williams construction allows at most one edge between vertex pairs (i, j) and it forbids overlaps in the initial sets; these two conditions are needed to handle maps having different contraction ratios $t_{i,j}$ on different edges, and are essential to obtain their formula for the Hausdorff dimension of the resulting set.

Theorem 2.5. *Let $\mathcal{G}$ be a connected labelled graph with vertices $V = \{1, 2, \dots, n\}$, and with edge label alphabet $\mathcal{A} = \{0, 1, 2, \dots, p-1\}$. Then there exist unique nonempty compact sets $\{K_i : i \in V\}$, each contained in $\mathbb{Z}_p$, that satisfy the set-valued functional relations*

$$(2.6) \quad K_i = \bigcup_{\{e: i(e)=i\}} \phi_e(K_{f(e)}), \quad i \in V.$$

Proof. This existence and uniqueness of the compact set-valued fixed point (2.6) follows from Hutchinson [12, Theorem 3.1]. The Hutchinson proof establishes that the K_v are obtained by the following iterative process. We start with initial sets $K_i^{(0)} = \mathbb{Z}_p$, and iteratively define

$$K_i^{(k+1)} := \bigcup_{\{e: i(e)=i\}} \phi_e(K_{f(e)}^{(k)}), \quad i \in V.$$

We have the sequence of closed sets

$$K_i^{(0)} \supseteq K_i^{(1)} \supseteq K_i^{(2)} \supseteq \dots,$$

and these converge downwards to the compact sets K_i , as $n \rightarrow \infty$. $\square$

Definition 2.6. Any set given as $Y := K_i$ for some sub-object K_i in the restricted p -adic graph directed system is called a *p -adic path set fractal*.

Many different construction pairs $(\mathcal{G}, i)$ can produce the same p -adic path set Y . One can use this freedom to make good choices for $\mathcal{G}$. For example, one may require $\mathcal{G}$ to be right-resolving, and in addition to have at most one directed edge between any directed pair of vertices, see the proof of Theorem 3.1(1) in Section 3.

2.3. Path sets. We give an alternate characterization of p -adic path set fractals in terms of the symbolic dynamics of their members. In [2] we introduced a notion of *path set* in abstract symbolic dynamics defined as follows. Let $X = \mathcal{A}^{\mathbb{N}}$ be the full one-sided shift space on $\mathcal{A}$. A *pointed graph* over an alphabet $\mathcal{A}$ consists of a pair $(\mathcal{G}, v)$, where $\mathcal{G} = (G, \mathcal{A})$ is a finite edge-labeled directed graph G , with edge labels drawn from an alphabet $\mathcal{A}$, and v a vertex of G . We let $V(\mathcal{G})$ and $\mathcal{E}(\mathcal{G})$ denote the set of vertices and directed edges of G , respectively. Following [2] we make a basic definition.

Definition 2.7. For a pointed graph $(\mathcal{G}, v)$ its associated *path set* (or *pointed follower set*) $X_{\mathcal{G}}(v) \subset \mathcal{A}^{\mathbb{N}}$ is the set of all infinite one-sided symbol sequences giving the successive labels of all one-sided infinite walks in $\mathcal{G}$ issuing from the distinguished vertex v . Many different $(\mathcal{G}, v)$ may give the same path set $\mathcal{P}$, and we call any such $(\mathcal{G}, v)$ a *presentation* of $\mathcal{P}$.

Now suppose $\mathcal{A} = \{0, 1, \dots, p-1\}$. Then the one-sided shift space $\mathcal{A}^{\mathbb{N}}$, topologized with the product topology, is homeomorphic to $\mathbb{Z}_p$ with its usual p -adic topology, where one identifies a symbol sequence $\alpha_0\alpha_1\alpha_2\cdots \in \mathcal{A}^{\mathbb{N}}$ with the p -adic expansion

$$x = \sum_{j=0}^{\infty} \alpha_j p^j = (\alpha_0, \alpha_1, \alpha_2, \dots)_p \in \mathbb{Z}_p.$$

We write $Y_{\mathcal{P}}$ to denote this path set viewed as a geometric object inside $\mathbb{Z}_p$, via the embedding map $j : \mathcal{A} \rightarrow \mathbb{Z}_p$.

The following result shows that this class of sets coincides with the *p -adic path set fractals* defined above.

Theorem 2.8. *Let $\mathcal{A} = \{0, 1, 2, \dots, p-1\}$ and Let $(\mathcal{G}, v)$ be path set data, with $\mathcal{G} = (G, V, E, \mathcal{L})$ being a labeled directed graph, with labels in $\mathcal{A} = \{0, 1, 2, \dots, p-1\}$. Let $Y_{\mathcal{P}} \subset \mathbb{Z}_p$ denote this path set, viewed as an embedded geometric object inside $\mathbb{Z}_p$. Then*

$$Y_{\mathcal{P}} = K_v,$$

where K_v is a p -adic path-set fractal on the same alphabet determined by the p -adic graph directed system with the same data $\mathcal{G} = (G, V, E, \mathcal{L})$. The converse also holds.

Proof. The correspondence proceeds by relating paths to the address labels of points in the fractal, compare Edgar [7, Sect. 4.3]. We study the set-valued iteration $K_i^{(k)}$ given in Theorem 2.5 for the p -adic path set fractal determined by $\mathcal{G} = (G, V, E, \mathcal{L})$. One proves by induction on $k \geq 0$ that for all vertices i that

$$K_i^{(k)} := \bigcup \left((\alpha_0 + \alpha_1 p + \dots + \alpha_{k-1} p^{k-1}) + p^k \mathbb{Z}_p \right)$$

where the set union is taken over label sequences $(\alpha_0, \dots, \alpha_{k-1})$ of legal walks in the directed graph $\mathcal{G}$ of length k starting from vertex i . The hypothesis that an exit edge exists from each vertex guarantees that all paths extend one step. The base case $k = 0$ holds since all $K_i^{(0)} = \mathbb{Z}_p$. For the induction step, we have $K_i^{(k+1)}$ is comprised of sets

$$\phi_e(K_{f(e)}^{(k)}) = j_e + \sum_{i=0}^{k-1} \alpha_i p^{i+1} + p^{k+1} \mathbb{Z}_p$$

where $(\alpha_0, \dots, \alpha_{k-1})$ are labels from a directed walk in $\mathcal{G}$ starting from vertex $f(e)$. But now $(j_e, \alpha_0, \dots, \alpha_{k-1})$ are vertices of a directed walk of length $k+1$ starting from vertex i , and all such walks are enumerated this way. This completes the induction step. Letting $k \rightarrow \infty$, for each vertex i these sets decrease to K_i , which is now identified with all infinite walks in $\mathcal{G}$ starting from vertex i . Choosing i to be the original marked vertex, we obtain $K_i = Y_{\mathcal{P}}$, as asserted. The converse is immediate. $\square$

In [2, Theorem 1.1] we observe that the class of path sets on a fixed alphabet is closed under finite unions and intersections. Now Theorem 2.8 implies that p -adic path set fractals inherit these closure properties.

2.4. p -adic path sets of cardinality one. We characterize path sets consisting of a single element; these give a dynamical characterization of the p -integral rational numbers $\mathbb{Q} \cap \mathbb{Z}_p$.

Theorem 2.9. (Single element p -adic path sets) *The p -adic path set fractals K in $\mathbb{Z}_p$ that consist of a single element are exactly those $K = \{r\}$ for which r is a p -integral rational number, i.e. $r \in \mathbb{Q} \cap \mathbb{Z}_p$.*

Proof. Given a path set $Y_{\mathcal{P}}$, given with a right-resolving presentation $\mathcal{P} = (\mathcal{G}, v_0)$, whose vertices are pruned to remove all vertices with no exit edges, can consist only of a single infinite path if there is exactly one exit edge from each vertex, and if the path is eventually periodic. The latter forces any element r to be a rational number in $\mathbb{Z}_p$. Conversely, we may easily construct a path set consisting of a single element giving the p -adic expansion of r . $\square$

3. HAUSDORFF DIMENSION OF p -ADIC PATH SETS

We relate p -adic path sets and graph-directed constructions on the real numbers, and use this to obtain a formula for their Hausdorff dimension. Recall that the *adjacency matrix* $A = A(G)$ of a directed graph G is a non-negative integer matrix whose rows and columns are numbered by the vertices of G (in the same order) with entry A_{ij} counting the number of directed edges outgoing from vertex i and incoming to vertex j .

Theorem 3.1. *Let p be a prime, and let X be a set of p -adic integers whose allowable p -adic expansions are described by the symbolic dynamics of a path set $\mathcal{P}$ on symbols $\mathcal{A} = \{0, 1, 2, \dots, p-1\}$.*

(1) *The map $\iota_p : \mathbb{Z}_p \rightarrow [0, 1] \subset \mathbb{R}$ taking $\alpha = \sum_{k=0}^{\infty} a_k p^k \in \mathbb{Z}_p$ to the corresponding real number with base p expansion*

$$\iota_p(\alpha) := \sum_{k=0}^{\infty} \frac{a_k}{p^{k+1}}$$

is a continuous map. Then the image set $K_0 := \iota_p(X) \subset [0, 1]$ is a construction sub-object K_v of a graph-directed fractal in the sense of Mauldin-Williams, whose edge similarity maps are all contracting similarity maps of $\mathbb{R}$ with contraction ratio $\frac{1}{p}$.

(2) *The Hausdorff dimensions of these sets are related by*

$$d_H(X) = d_H(K_0).$$

(3) *If $(\mathcal{G}, v_0)$ is a right-resolving presentation for the path set X , then*

$$(3.1) \quad d_H(X) = d_H(K_0) = \log_p \alpha,$$

where $\alpha = \sigma(A(\mathcal{G}))$ is the spectral radius of the adjacency matrix $A = A(\mathcal{G})$ of the underlying directed graph $\mathcal{G}$ of $\mathcal{G}$.

Remark 3.2. (1) We observed in Theorem 1.9 of [2] that the topological entropy of a path set X , with a right-resolving presentation is given by

$$h_{top}(X) = \log \alpha,$$

where α is the spectral radius as above. We deduce that for a p -adic path set fractal, its Hausdorff dimension is

$$d_H(X) = \frac{h_{top}(X)}{\log p}.$$

This extends to path sets a well-known result in the shift-invariant case (Furstenberg [11, Prop. III.1]).

(2) The right-resolving property assumed in (3) is needed to guarantee equality of the Hausdorff dimension with $\log_p \alpha$. For a general presentations $(\mathcal{G}, v_0)$ of X the adjacency matrix counts the growth rate of paths, which upper bounds the number of distinct sequences of path labels. That is, one always has

$$d_H(X) \leq \log_p \sigma(A(\mathcal{G})).$$

(3) The allowable class of values α that may occur are the special class of positive real algebraic integers called *Perron numbers*, that were introduced by Lind [16].

Proof. (1) We first find a good presentation $(\mathcal{G}, v)$ of the path set $X = X_{\mathcal{P}}$. A presentation is *right-resolving* if for each vertex v of $\mathcal{G}$, all exit edges have different labels. By a standard construction, a path set always has a right-resolving presentation in which $\mathcal{G}$ is connected and every vertex has an exit edge ([2, Theorem 3.2], compare [17, Theorem 3.3.2]). We call a presentation *right-separating* if the underlying directed graph G of $\mathcal{G}$ has no multiple edges.

Claim. *There exists a presentation $(\mathcal{G}, v)$ of X in which $\mathcal{G}$ is both right-resolving and right-separating.*

To show the claim, given a right-resolving presentation $(\mathcal{G}', v')$, we show it may be converted to a right-separating presentation by making use of a vertex-splitting construction, as follows. Suppose that a vertex v of $\mathcal{G}'$ has $k \geq 2$ labelled edges from a vertex w , which necessarily has distinct labels. We create a new labelled graph $\mathcal{G}''$ that retains all vertices of $\mathcal{G}$ except v and replaces v by k vertices $v^{(i)}$, $1 \leq i \leq k$.

Case 1. $w \neq v$.

In this case we assign a single new labelled edge from w to each $v^{(i)}$ such that as i varies, the corresponding edge labels exhaust the k labels of edges from w to v . The exit edges assigned each $v^{(i)}$ each duplicate the exit edges from v , both in multiplicity and in labels, with self-loops of v corresponding to self-loops of $v^{(i)}$. The entering edges to $v^{(1)}$ will all be the same in multiplicity and in labels as for v , while for $v^{(i)}$ with $i \geq 2$ there will be no entering edges from the rest of the graph, with the exception of self-loops, assigned as above. Finally all edges between any two vertices distinct from the $v^{(i)}$ will be the same as in the original graph.

Case 2. $w = v$

In this case v has k self-loops, all with different labels since the right-resolving property is assumed. In this case, by identifying the k loop labels with $\{1, 2, \dots, k\}$ in some fixed fashion, we assign an directed edge from $v^{(i)}$ to $v^{(j)}$ with edge label corresponding to $i + j \pmod k$, for $1 \leq i, j \leq k$. The other exit edges assigned each $v^{(i)}$ duplicate the exit edges of v , in multiplicity and labels (excluding self-loops). The entering edges of $v^{(1)}$ will all the same in multiplicity and labels as for v (excluding self-loops). All other $v^{(i)}$ have no entering edges at all from other vertices of the original graph $v' \neq v$.

We next assert that the path sets from all states $(\mathcal{G}'', w')$ for $w' \neq v^{(i)}$ agree with those of $(\mathcal{G}', w')$, while all path sets $(\mathcal{G}'', v^{(i)})$ agree identically with that of $(\mathcal{G}', v)$. This assertion may be established by viewing $\mathcal{G}'$ as a covering of $\mathcal{G}$ which preserves edge labels, which has the k vertices $v^{(i)}$ lying above vertex v , and all other vertices agreeing. One may check that each edge in $\mathcal{G}'$ from a given initial vertex v' lifts uniquely to a suitable vertex and edge above it (here $v' = w$ is the only interesting case), except that self-loops from v lift to a self-loop for any initial vertex $v^{(i)}$. After the first step, any path lifts uniquely. Conversely any labelled path in the lifted graph projects to an allowable labelled path in $\mathcal{G}'$. The assertion follows.

This construction has the feature that the new graph $\mathcal{G}''$ is still right-resolving, so it may be repeated. In doing so, we must eventually arrive at a right-resolving and right-separating presentation. Assign to each vertex an integer invariant that is the product of the multiplicities of all entering edges. When a vertex is split, this invariant decreases for all of the k descendants $v^{(i)}$, and remains the same for all

other vertices of the graph. By the well-ordering of $\mathbb{N}$, the splitting procedure will eventually halt at a right-separating presentation. This establishes the claim.

We now suppose given a right-resolving and right-separating presentation $(\mathcal{G}, v)$ of X , and we proceed to map this to a corresponding graph-directed construction. We arrange to shift the image sets by integers in such a way as to satisfy the non-overlapping condition in the Mauldin-Williams construction. View the vertices of $\mathcal{G}$ as labelled with integers $0 \leq i \leq n = |V(\mathcal{G})| - 1$, with the We let X_j denote the path set associated to $(\mathcal{G}, j)$, with $X_0 = X$. We map the set X_j under the map $\iota_p + 2j$ to the image set

$$K_j := \iota(X_j) + 2j \subset [2j, 2j + 1],$$

with the integer shifts making all sets K_j disjoint in $\mathbb{R}$.

We now define a Mauldin-Williams geometric graph-directed construction in $\mathbb{R}$, with all image sets contained in the compact set $[0, 2n]$, for which the sets $(K_0, K_1, \dots, K_n)$ form the construction sub-objects. The underlying graph G of $\mathcal{G}$ above has the correct properties to be a graph in the Mauldin-Williams construction: it is connected, has at most one directed edge between any ordered pair of vertices, and each vertex has at least one exit edge. To each directed labelled edge $e = [i_1, i_2]$ of $\mathcal{G}$ with label j we associate real-valued map

$$\phi_e(x) = \frac{(x - 2i_2) + j}{p} + 2i_1.$$

By the basic theorem of Hutchinson [12, Theorem 3.1], this construction has a unique compact attractor consisting of a collection of compact sets $\{K_j : 0 \leq j \leq n\}$. These sets K_j are directly by the Mauldin-Williams (iterative) geometric graph-directed construction using [21, Theorem 1] starting with the (disjoint) initial sets $J_j := [2j, 2j + 1]$ ($v \in V(\mathcal{G})$). In particular, the non-overlapping property (a) of all $T_{i,j}(J_j)$ follows from the right-resolving property of the graph $\mathcal{G}$. These sets are compact, with non-empty interior, and after m -iterations, we have sets $J_v^{(m)}$ ($v \in V(\mathcal{G})$) which form nested sequences, and we then obtain $K_v := \bigcap_m J_v^{(m)}$. The Mauldin-Williams construction object is $K = \bigcup_{v \in V(\mathcal{G})} K_v$. The set $K_i \subset [2i, 2i + 1]$, and $K_0 = \iota(X)$ since $K_0 \subset [0, 1]$ and the underlying symbol sequences of X and of K_0 agree. (Some points in K_0 have two preimages under the map ι_p , and only one of the two preimage symbol sequences occur in this case.)

(2) The definition of p -adic Hausdorff dimension is quite similar to Hausdorff dimension for real numbers on the interval $[0, 1]$, cf. Abercrombie [1]. An ϵ -coverings of X is a covering of X by a countable collection of p -adic open balls all having diameter at most ϵ , and considers the quantities

$$m_\beta(X) := \lim_{\epsilon \rightarrow 0} \inf_{\epsilon\text{-cover}} (\text{Vol}(B(x_i, \epsilon_i)))^\beta,$$

in which the data $\{(x_i, \epsilon_i) : i \geq 1\}$ describes the covering, specifying center x_i and radius ϵ_i of p -adic disks, and with $0 < \epsilon_i \leq \epsilon$, and $\text{Vol}(S)$ denotes the usual p -adic measure of $S \subset \mathbb{Z}_p$. There is a cutoff value β_0 such that $m_\beta(X) = \infty$ for $\beta > \beta_0$ and $m_\beta(X) = 0$ for $\beta < \beta_0$; this is the Hausdorff dimension of X . We use the following basic fact, following [15, Sec. 3.2].

Claim. *There is a continuous (and almost one-to-one) mapping $\iota_p : \mathbb{Z}_p \rightarrow [0, 1]$ which sends a p -adic number $\lambda = (\dots d_2 d_1 d_0)_p$ to the real number with base p expansion $.d_0 d_1 d_2 \dots$. This mapping preserves Hausdorff dimension of sets, i.e a p -adic set X and its image $\iota_p(X)$ have the same Hausdorff dimension.*

To verify the claim, note that one can expand each set in a p -adic covering of a set X to a closed-open disk

$$B(m, p^j) = \{x \in \mathbb{Z}_p : x \equiv m \pmod{p^j}\}$$

(which has diameter $\frac{1}{p^j}$), with at most a factor of p increase in diameter, and similarly one can inflate any real covering to a covering with ternary intervals $[\frac{m}{p^j}, \frac{m+1}{p^j}]$ with at most a factor of p increase in diameter. But these special intervals are assigned the same diameter under their respective metrics, and this can be used to show the Hausdorff dimensions of X and $\iota(X)$ coincide. (The Hausdorff measures of the resulting sets is not proved to coincide by this argument.)

The truth of the claim immediately yields $d_H(K_0) \equiv d_H(\iota(X)) = d_H(X)$, as asserted.

(3) Suppose we are given a right-resolving presentation $(\mathcal{G}, v_0)$ of the p -adic path set. Assume first that this presentation is right-separating. In that case we can directly make use of Proposition 2.3 of Mauldin-Williams. The set K_{v_0} is connected by paths to every vertex of the graph $\mathcal{G}$, so that the vertex set $C_1 := S(\mathcal{G})$ reaches all strongly connected components of $\mathcal{G}$. By Proposition 2.3 we obtain that the Hausdorff dimension of K_{v_0} is the same as that of the full construction object $K = \bigcup_v K_v$.

The Hausdorff dimension of K_v is computed in [21] in terms of the adjacency matrix of a graph encoding the contraction maps. In our case all nonzero maps for $\mathcal{G}^*$ are similarities with constant ratio $\frac{1}{p}$, which yields the matrix

$$A_\beta = \left(\frac{1}{p}\right)^\beta A(\mathcal{G}^*), = \left(\frac{1}{p}\right)^\beta A(\mathcal{G})^T$$

where $A(\mathcal{G})$ is the adjacency matrix for the underlying directed graph G , and $A(\mathcal{G})^T$ is its transpose. Here $A(\mathcal{G})_{v,w}$ counts the number of directed edges from vertex v to vertex w . Now we assume the map $\mathcal{G}$ is right-resolving (this guarantees a certain non-overlap condition for addresses in the final sets K_v). We set

$$\Phi(\beta) := \text{Spectral radius of } A_\beta := \lambda_\beta.$$

Now the special form of A_β yields

$$\Phi(\beta) = \lambda_\beta = \lambda_0 p^{-\beta},$$

in which λ_0 is the spectral radius of $\sigma(A(\mathcal{G}))$, which equals $\sigma(A(\mathcal{G})^T)$. By [21, Theorem 3] the full construction object has Hausdorff dimension

$$d_H(K) = \alpha$$

where $\Phi(\alpha) = 1$. This gives

$$\alpha = \log_p \lambda_0 = \log_p \sigma(A[\mathcal{G}]),$$

as required.

It remains to treat the general case, in which $(\mathcal{G}, v_0)$ is not necessarily right-separating, and show that the formula for Hausdorff dimension continues to hold. To handle this case, we study the effect of the state-splitting construction introduced earlier to convert a right-resolving presentation $(\mathcal{G}', v')$ of a path set to one $(\mathcal{G}, v_0)$ that is also right-separating. It suffices to show that every step of this procedure preserves the value of the spectral radius (Perron eigenvalue) of the associated nonnegative matrix $[A(\mathcal{G}'')]$. If this is shown, then the already proved formula for the Hausdorff dimension for the p -adic path set associated to $(\mathcal{G}, v_0)$ will carry over

to that for $(\mathcal{G}', v')$. We check the spectral radius is preserved. We use the known fact that the spectral radius of a nonnegative matrix A is given by

$$\sigma(A) = \lim_{n \rightarrow \infty} (N_n(A))^{\frac{1}{n}},$$

in which $N_n(A) = \mathbf{e}^T A^n \mathbf{e}$, where $\mathbf{e} = [1, 1, \dots, 1]^T$ is a column vector. Here $N_n(A)$ counts the number of directed paths of length n between all pairs of vertices of A . (The existence of the limit is part of the assertion.) We use the fact that $\mathcal{G}''$ is a covering of $\mathcal{G}'$ and that all (labelled) paths of $\mathcal{G}'$ lift uniquely to paths of $\mathcal{G}''$, with the exception of paths that have starting vertex v , which have k distinct lifts, where k was the number of vertices of $\mathcal{G}$ that were split. From this we conclude that

$$N_n(A(\mathcal{G})) \leq N_n(A(\mathcal{G}'')) \leq k N_n(A(\mathcal{G})),$$

and since k is constant, we conclude that

$$\sigma(A(\mathcal{G}')) \leq \sigma(A(\mathcal{G}'')) \leq \lim_{n \rightarrow \infty} k^{\frac{1}{n}} N_n(A(\mathcal{G}'))^{\frac{1}{n}} = \sigma(A(\mathcal{G}')),$$

giving the result. $\square$

Proof of Theorem 1.1. This is immediate from Proposition 3.1. It represents the set X_{v_0} as a pullback from one set in a graph-directed fractal of the type of Mauldin and Williams[21], which has identical symbolic dynamics, with the Mauldin-Williams non-overlapping condition enforced. $\square$

4. ADDITION OF p -ADIC PATH SET FRACTALS

We analyze the effect on p -adic path sets of addition with rational numbers $r \in \mathbb{Q}$, viewing $\mathbb{Q}$ as a subfield of $\mathbb{Q}_p$. We describe algorithms which when given a realization $(\mathcal{G}, v_0)$ of a path set X , will produce a realization of $X + r$, sometimes under restrictions on X .

4.1. Sum of a path set and a p -integral rational number. Theorem 1.2 is an immediate corollary of the following stronger result. Recall that a p -integral rational number r is any $r \in \mathbb{Q} \cap \mathbb{Z}_p$.

Theorem 4.1. *Let $\mathcal{P}$ be a path set having a right-resolving presentation $(\mathcal{G}, v_0)$ having V vertices. Suppose that r is a p -integral rational number, which has a p -adic expansion with pre-periodic part of length Q_0 and a periodic part of period Q . Then the additively shifted set $\mathcal{P}' := \mathcal{P} + r$ is a p -adic path set, which has a right-resolving presentation having at most $2p(Q_0 + Q)V$ vertices.*

Proof. We give an explicit construction, starting from the given right-resolving presentation $(\mathcal{G}, v_0)$.

We suppose first that r has a purely periodic p -adic expansion $(c_0 c_1 c_2 \dots)_p$, of period Q , with $c_{j+Q} = c_j$, and write

$$r = \sum_{j=0}^{\infty} c_j p^j = \left(\sum_{j=0}^{Q-1} c_j p^j \right) \left(\sum_{k=0}^{\infty} p^{kQ} \right).$$

We aim to construct a right-resolving presentation $(\mathcal{G}', \mathbf{w}_0)$ for $\mathcal{P}' := \mathcal{P} + r$. The vertex states of $\mathcal{G}'$ will be labelled $\mathbf{w} = (v, f, e, a)$, in which:

- (i) v denotes a vertex of $\mathcal{G}$;
- (ii) f with $0 \leq f \leq Q-1$ denotes a place-marker in the periodic portion of the p -adic expansion of r ,

- (iii) e keeps track of the current amount of “carry-digit” information not yet incorporated in the sum-set p -adic expansion,
- (iv) a with $0 \leq a \leq p-1$ denotes an edge label value.

The initial vertex will be $\mathbf{w}_0 = (v_0, 0, 0, 0)$. We will establish an upper bound on the maximum size $e \leq 2$ in the analysis below.

The exit edges of $\mathcal{G}'$ map a vertex $\mathbf{w}$ to $\mathbf{w}' = (v', f', e', a')$ in which there is a directed labelled edge $(v, v') \in \mathcal{G}$, with label ℓ_1 satisfying $\ell_1 = a'$, and the value $f' \equiv f + 1 \pmod{Q}$. The edge label ℓ' assigned to this edge will be $0 \leq \ell \leq p-1$ with

$$(4.1) \quad \ell' \equiv e + \ell_1 + c_f \pmod{p-1},$$

and the value e' is required to satisfy

$$(4.2) \quad e' = \frac{1}{p}(e + \ell_1 + c_f - \ell').$$

We define the graph $\mathcal{G}'$ to consist of all states reachable from the initial vertex $\mathbf{w}_0$. We first show that all reachable vertices satisfy the bound $e \leq 2$; this shows that the graph $\mathcal{G}'$ is finite and also bounds its size. The bound is proved by induction on the number of steps n along a directed path. The base case $n = 0$ has $e = 0$. For the induction step, using the rule above

$$e' = \frac{1}{p}(e + \ell_1 + c_f - \ell') \leq \frac{1}{p}(2 + (p-1) + (p-1)) \leq 2,$$

completing the induction step.

We next show that the presentation $(\mathcal{G}', \mathbf{w}_0)$ has path set $\mathcal{P}'$ satisfying $\mathcal{P}' = \mathcal{P} + r$. We first check that $\mathcal{G}'$ is right-resolving. To see this, note that the exit edges from a vertex $\mathbf{w}$ correspond to exit edges from vertex v in the right-resolving graph $\mathcal{G}$, whence any two edges have different values of ℓ_1 . Now the exit edge label ℓ' is an invertible linear function of ℓ_1 by (4.4), since the values e and c_f are fixed by $\mathbf{w}$, so all exit edges have distinct labels, as asserted.

We next observe that a lifted path in $\mathcal{G}'$ uniquely determines an underlying path in $\mathcal{G}$. This follows since the path label value ℓ_1 is uniquely recoverable from the path label ℓ' and the vertex data on $\mathcal{G}$, using (4.5), since e' is known and c_f is known from the vertex label f . The underlying path on $\mathcal{G}$ determines

$$x_n = \sum_{k=0}^{n-1} a_k p^k, \quad i = 1, 2,$$

corresponding to the initial part of the p -adic expansion of a value $x \in \mathcal{P}$ being determined by the $\mathcal{G}$ -path. (The values a_k are the successive labels ℓ_1 along the $\mathcal{G}$ -path.) Conversely, each path in $\mathcal{G}$ with initial vertex v_0 lifts to a unique path in $\mathcal{G}'$ with initial vertex $\mathbf{w}_0$. Given a vertex $\mathbf{w}$, a labelled edge (v, v') with label ℓ_1 determines the values $a' = \ell_1$ and e' a unique vertex $\mathbf{w}'$ that $\mathbf{w}$ connects to.

We now show that for a path in $\mathcal{G}$ determining $x \in \mathcal{P}$, the labels of the lifted path in $\mathcal{G}'$ suffice to compute the value $x + h \in \mathcal{P}'$. We prove this by induction on n , for the n -step initial path. The successive edge labels $\{\ell'_i : 0 \leq i \leq n-1\}$ of the lifted path in $\mathcal{G}'$ with the end vertex data $e' = e_n$ determine the quantity

$$y_n := \sum_{k=0}^{n-1} \ell'_k p^k + e_n p^n.$$

We establish by induction that

$$(4.3) \quad y_{n-1} = x_{n-1} + r_{n-1},$$

in which

$$r_m := \sum_{j=0}^{m-1} c_j p^j,$$

is a truncated version of the p -adic expansion of r . By the induction hypothesis,

$$\begin{aligned} x_n + r_n &= (x_{n-1} + r_{n-1}) + a_{n-1}p^{n-1} + c_{n-1}p^{n-1} \\ &= \left(\sum_{k=0}^{n-2} \ell'_k p^k + e_{n-1}p^{n-1} \right) + a_{n-1}p^{n-1} + c_{n-1}p^{n-1} \\ &= \left(\sum_{k=0}^{n-1} \ell'_k p^k \right) + e_n p^n, \end{aligned}$$

the last equality holding by virtue of (4.5), using $c_f = c_{n-1}$ and $\ell_1 = a_n$. This completes the induction step, proving (4.3).

Now the lifted path data yields the p -adic limit

$$\lim_{n \rightarrow \infty} \sum_{k=0}^{n-1} \ell'_k p^k = \lim_{n \rightarrow \infty} (x_n + r_n) - \lim_{n \rightarrow \infty} e_n p^n = x + r.$$

We conclude that the lifted path of $\mathcal{G}'$ corresponding to $x \in \mathcal{P}$ determines the point $x + r \in \mathcal{P}'$. It follows that $\mathcal{P}' = \mathcal{P} + r$, as asserted. This completes the argument for r having a purely periodic p -adic expansion.

It remains to extend this construction to the case where r has a preperiodic part of length Q_0 , and to upper bound the number of states in the constructed presentation $(\mathcal{G}, \mathbf{w})$. The extension is routine: we add extra vertices $\mathbf{w} := (v, d_j, e, a)$ to $\mathcal{G}$, in which v denotes a vertex of $\mathcal{G}$, d_j marks the j -th preperiodic digit of r , $1 \leq j \leq Q_0$. The exit edges of $\mathcal{G}'$ map a vertex to $\mathbf{w}' = (v', d_{j+1}, e', a')$ in which there is a directed labelled edge $(v, v') \in \mathcal{G}$, with label ℓ_1 satisfying $\ell_1 = a'$. The edge label ℓ' assigned to this edge will be $0 \leq \ell \leq p-1$ with

$$(4.4) \quad \ell' \equiv e + \ell_1 + d_j \pmod{p-1},$$

and the value e' is required to satisfy

$$(4.5) \quad e' = \frac{1}{p}(e + \ell_1 + d_j - \ell').$$

The final preperiodic digit exit edges go to vertices $\mathbf{w} := (v', f_0, e', a')$ in the earlier set.

It is straightforward to check that the underlying labelled graph has the right-resolving property. Finally one must check this extension preserves the lifting property of paths, we omit details.

It remains to upper bound the total number of vertices in the graph $(\mathcal{G}, \mathbf{w})$. The preperiodic part contributes at most $2Q_0 V p$ vertices, and the periodic part contributes at most $2Q V p$ vertices. $\square$

Remark 4.2. The key features in the proof are: (i) the p -adic carry digits propagate to higher powers of p and do not disturb earlier p -adic digits; (ii) the size of the

carry digits is bounded above. Here property (i) fails in the real number case, and there is no real number analogue of this result.

4.2. Set sum of two p -adic path sets. We show that the set sum of two p -adic path sets is itself a path set, establishing Theorem 1.3. This proof is constructive, but it no longer produces a right-resolving presentation.

Proof of Theorem 1.3. We suppose that $\mathcal{P}_1$ and $\mathcal{P}_2$ have associated right-resolving pointed graph presentations $(\mathcal{G}_i, v_{0,i})$, $(i = 1, 2)$. We use these presentations to directly construct a presentation $(\mathcal{G}_{1,2}, \mathbf{w}_0)$ for $\mathcal{P}_1 + \mathcal{P}_2$, certifying that it is a path set.

To begin the construction, $\mathcal{G}_{1,2}^+$ will have vertices labelled $\mathbf{w} := (v_{j,1}, v_{k,2}, e, a)$ where $v_{j,1} \in V(\mathcal{G}_1)$, $v_{k,2} \in V(\mathcal{G}_2)$, and $e \geq 0$ is an integer encoding “carry-digit” information, and $0 \leq a \leq p-1$ specifies an allowed edge entry label in $\mathcal{G}_1$. A bound on the maximum size of e will be established below.

The exit edges from vertex $\mathbf{w}$ go to a new vertex $\mathbf{w}' = (v_{j',1}, v_{k',2}, e', a')$ in which

(a) there is a directed edge of $\mathcal{G}_1$ from $v_{j,1}$ to $v_{j',1}$ having label ℓ_1 satisfying $\ell_1 = a'$;

(b) a directed edge of $\mathcal{G}_2$ from $v_{k,2}$ to $v_{k',2}$ with label ℓ_2 , and

(c) the constructed edge is assigned the label ℓ , $0 \leq \ell \leq p-1$, determined by

$$\ell \equiv e + \ell_1 + \ell_2 \pmod{p}, \quad 0 \leq \ell \leq p-1$$

(d) the new carry-digit is

$$e' = \frac{1}{p}(e + \ell_1 + \ell_2 - \ell) \geq 0.$$

The initial pointed vertex of the graph $\mathcal{G}_{1,2}^+$ is $\mathbf{w}_0 := (v_{0,1}, v_{0,2}, 0, 0)$. We now define $\mathcal{G}_{1,2}^+$ to consist of all vertices above reachable by a directed path starting from $\mathbf{w}_0$. We show this is a finite graph by establishing that the carry digit in any reachable vertex satisfies $e \leq 2$. This follows by induction on the length of the path. The base case is $n = 0$ where it holds vacuously. For the induction step, we upper bound the new value of e via

$$e' = \frac{1}{p}(e + \ell_1 + \ell_2 - \ell) \leq \frac{1}{p}(2 + (p-1) + (p-1)) \leq 2,$$

completing the induction step.

To see that the path set produced by this graph is indeed $\mathcal{P}_1 + \mathcal{P}_2$, we prove by induction on $n \geq 0$ that being at a vertex $\mathbf{w}$ at step n says that the first n p -adic digit symbols of an n -step path $x_1 + x_2$ have been computed, namely we have

$$x_{i,n} = \sum_{k=0}^{n-1} a_{k,i} p^k, \quad i = 1, 2,$$

are such that

$$x_{1,n} + x_{2,n} = \sum_{k=0}^{n-1} b_k p^k + e p^n,$$

where $e = e_n$ is the current carry-digit, and the b_i are the edge labels produced so far in the graph $\mathcal{G}_{1,2}^+$. Suppose the next directed edge moves to a vertex $\mathbf{w}' = \mathbf{w}_{n+1}$, with data (e', a') . Then we have $a' = \ell_1 = a_{n,1}$ and

$$e' = \frac{1}{p}(\ell_1 + \ell_2 - \ell + e) = \frac{1}{p}(a + \ell_2 - \ell + e)$$

Since e, a, ℓ are all known, this equation uniquely determines the label $\ell_2 = a_{n,2}$. Since both $\mathcal{G}_1$ and $\mathcal{G}_2$ are right-resolving, the edges (j, j') and (k, k') with the labels $a_{n,1}, a_{n,2}$ are legal steps uniquely determines the edges updating $x_{1,n}, x_{2,n}$ to $x_{1,n+1}, x_{2,n+1}$. Now the definition of edge labels in $\mathcal{G}_{1,2}^+$ assigns the label $b_n := \ell$ to the edge of $\mathcal{G}_{1,2}^+$ and $e' = e_{n+1}$ in

$$x_{1,n+1} + x_{2,n+1} = \sum_{k=0}^n b_k p^k + e' p^n,$$

completing the induction step. $\square$

Remark 4.3. The path set presentation $(\mathcal{G}_{1,2}^+, \mathbf{w}_0)$ for $\mathcal{P}_1 + \mathcal{P}_2$ given in this construction is generally far from right-resolving. A geometric interpretation is that this occurs because some values $x = x_1 + x_2 \in \mathcal{P}_1 + \mathcal{P}_2$ may have more than one representation (x_1, x_2) . This construction produces a separate path for each pair (x_1, x_2) , so more than one path can yield the same sequence of labels.

5. MULTIPLICATION AND p -ADIC PATH SETS

5.1. Multiplication by rational numbers. We give constructive proof for multiplication by rational numbers of specific types.

Theorem 5.1. *Let $\mathcal{P}$ be a path set having a right-resolving presentation $(\mathcal{G}, v_0)$ having V vertices. Let $M \geq 2$ be a positive integer with $\gcd(p, M) = 1$.*

- (1) *For $r = M$ the multiplicatively shifted set $\mathcal{P}' := M\mathcal{P}$ is a p -adic path set which has a right-resolving presentation having at most $(M+1)V$ vertices.*
- (2) *For $r = \frac{1}{M}$ the multiplicatively shifted set $\mathcal{P}' := \frac{1}{M}\mathcal{P}$ is a p -adic path set which has a right-resolving presentation having at most $(M+1)V$ vertices.*
- (3) *For $r = -1$ the multiplicatively shifted set $\mathcal{P}' := -\mathcal{P}$ is a p -adic path set which has a right-resolving presentation having at most $(p+1)V$ vertices.*
- (4) *For $r = p^k$, with integer $k \geq 0$, the multiplicatively shifted set $\mathcal{P}' := p^k\mathcal{P}$ is a p -adic path set which has a right-resolving presentation having $k+V$ vertices.*

Proof. Given right-resolving presentation $(\mathcal{G}, v_0)$ of $\mathcal{P}$ with V states, we give an explicit construction of a right-resolving presentation $(\mathcal{G}', \mathbf{w}_0)$ for $r\mathcal{P}$, for each given r . The constructions in cases (1)-(3) are similar.

(1) We have $p \nmid M$, and construct a presentation $(\mathcal{G}', \mathbf{w}_0)$ for $M\mathcal{P}$. The vertices of $\mathcal{G}'$ will be pairs $\mathbf{w} = (v, c)$, in which v is a vertex of $\mathcal{G}$, c is a carry digit, which will satisfy $0 \leq c \leq M$. The initial vertex is $\mathbf{w}_0 := (v_0, 0)$. The exit edges from a vertex $\mathbf{w}$ to a vertex $\mathbf{w}' = (v', c')$ will occur only if there is at least one edge from v to v' . Given such an edge of $\mathcal{G}$ with label ℓ , we assign a corresponding edge of $\mathcal{G}'$ with label ℓ' given by

$$(5.1) \quad \ell' = M\ell + c \pmod{p}, \quad 0 \leq \ell' \leq p-1,$$

which is well-defined since $(p, M) = 1$. We require that the new carry digit be

$$(5.2) \quad c' := \frac{1}{p}(c + M\ell - \ell').$$

We define $(\mathcal{G}', \mathbf{w}_0)$ to the graph obtained taking all vertices reachable from $\mathbf{w}_0$ in the above construction. We prove that all reachable states have $0 \leq c \leq M$ by induction on the number of steps n on a minimal path to such a state. The base

case $n = 0$ is true, since $c = 0$, and the induction step follows by observing from (5.2) that

$$c' \leq \frac{1}{p}(M + M(p-1)) \leq M.$$

We conclude that the graph $\mathcal{G}'$ has at most $(M+1)V$ states.

The pointed graph $(\mathcal{G}', \mathbf{w}_0)$ now determines a path set $\mathcal{P}'$. We first observe that $\mathcal{G}'$ is right-resolving. We argue by contradiction. If not, there would be two exit edges of some vertex $\mathbf{w} = (v, c)$ of $\mathcal{G}$ having the same value of ℓ' . But then by (5.1) the underlying edges of $(\mathcal{G}, v_0)$ would have the same value of ℓ , contradicting the right-resolving property of $(\mathcal{G}, v_0)$.

It remains to show that if $\mathcal{P}'$ is the resulting path set, then $\mathcal{P}' = M\mathcal{P}$. Consider an infinite path in $\mathcal{G}$, with image

$$x_\infty = \sum_{j=0}^{\infty} \ell_j p^j \in \mathcal{P}$$

We assert the corresponding output path

$$y_\infty = \sum_{j=1}^{\infty} \ell'_j p^j \in \mathcal{P}'$$

has

$$y_\infty = \frac{1}{M} x_\infty.$$

We verify this by an induction on the length of a finite path approximating x_∞ . Let $v_0, v_1, \dots, v_n$ be states on a path in sG with edge labels $\ell_0, \ell_1, \ell_{n-1}$. Associated to this path is

$$x_n = \ell_0 + \ell_1 p + \dots + \ell_{n-1} p^{n-1}.$$

By construction we obtain

$$y_n = \sum_{j=0}^{n-1} \ell'_j p^j.$$

Now we prove by induction on $n \geq 1$ that

$$y_n = Mx_n + e_n p^n$$

where $e_n = c'$, where c' is the carry value at the final vertex v_n . The base case $n = 1$ is clear, and for the induction step ($e_n = c, e_{n+1} = c'$) we get, using (5.2),

$$Mx_{n+1} = M(x_n + \ell_n p^n) = y_n + (e_n + M\ell_n)p^n = y_n + e_{n+1}p^{n+1}$$

Now we use $|e_n p^n|_p \rightarrow 0$ as $n \rightarrow \infty$, whence $My_\infty = x_\infty$ establishing the result.

(2) We have $p \nmid M$. The vertices of $\mathcal{G}_M$ will be pairs $\mathbf{w} = (v, c)$, in which v is a vertex of $\mathcal{G}$, and $c \geq 0$ is a carry digit, initially unbounded, which we later show will satisfy $0 \leq c \leq M$. The initial vertex is $bw_0 := (v_0, 0)$. The exit edges from a vertex $\mathbf{w}$ to a vertex $\mathbf{w}' = (v', c')$ will occur only if there is at least one edge in $\mathcal{G}$ from v to v' . Given such an edge of $\mathcal{G}$ with label ℓ , we assign a corresponding edge of $\mathcal{G}'$ with label ℓ' given by

$$(5.3) \quad M\ell' = \ell - c \pmod{p}, \quad 0 \leq \ell' \leq p-1,$$

which is well-defined since $(p, M) = 1$. We require that the new carry digit be

$$(5.4) \quad c' := \frac{1}{p}(c + M\ell' - \ell).$$

We define $(\mathcal{G}', \mathbf{w}_0)$ to be the graph obtained by including only the vertices reachable from $\mathbf{w}_0$ in the above construction. We prove that all reachable states have $0 \leq c \leq M$ by induction on the number of steps n on a minimal path to such a state. The base case $n = 0$ is true, since $c = 0$, and the induction step follows by observing from (5.4) that

$$c' \leq \frac{1}{p}(M + M(p-1)) \leq M.$$

We conclude that the graph $\mathcal{G}'$ has at most $(M+1)V$ states.

The pointed graph $(\mathcal{G}', \mathbf{w}_0)$ now determines a path set $\mathcal{P}'$. We first observe that $\mathcal{G}'$ is right-resolving. We argue by contradiction. If not, there would be two exit edges of some vertex $\mathbf{w} = (v, c)$ of $\mathcal{G}$ having the same value of ℓ' . But then by (5.3) the underlying edges of $(\mathcal{G}, v_0)$ would have the same value of ℓ , contradicting the right-resolving property of $(\mathcal{G}, v_0)$.

We now verify that the path set $\mathcal{P}'$ equals $\frac{1}{M}\mathcal{P}$, still supposing $(p, M) = 1$. Consider an infinite path in $\mathcal{G}$, with image

$$x_\infty = \sum_{j=0}^{\infty} \ell_j p^j \in \mathcal{P}$$

We assert the corresponding output path

$$y_\infty = \sum_{j=1}^{\infty} \ell'_j p^j \in \mathcal{P}'$$

has

$$y_\infty = \frac{1}{M}x_\infty.$$

We verify this by induction on finite paths. Let $v_0, v_1, \dots, v_n$ be states on a path in sG with edge labels $\ell_0, \ell_1, \ell_{n-1}$. Associated to this path is

$$x_n = \ell_0 + \ell_1 p + \dots + \ell_{n-1} p^{n-1}.$$

By construction we obtain

$$y_n = \sum_{j=0}^{n-1} \ell'_j p^j.$$

Now we prove by induction that

$$My_n = x_n + e_n p^n$$

where $e_n = c'$ the carry value at the final vertex v_n . For the induction step ($e_n = c, e_{n+1} = c'$) we get, using (5.4)

$$My_{n+1} = M(y_n + \ell'_n p^n) = x_n + (e_n + M\ell'_n)p^n = x_{n+1} + e_{n+1}p^{n+1}$$

Now we use $|e_n p^n|_p \rightarrow 0$ as $n \rightarrow \infty$, whence $My_\infty = x_\infty$ establishing the result.

(3) We construct a right-resolving presentation $(\mathcal{G}'', \mathbf{w}_0)$ for $(-1)\mathcal{P}$, as follows.

The vertices of $\mathcal{G}'$ will be pairs $\mathbf{w} = (v, c)$, in which v is a vertex of $\mathcal{G}$, and c is a carry digit, which may take values 0 or -1 . The initial vertex will be $bw_0 := (v_0, 0)$. The exit edges from a vertex $\mathbf{w}$ to a vertex $\mathbf{w}' = (v', c')$ will occur only if there is at least one edge in $\mathcal{G}$ from v to v' . Given such an edge of $\mathcal{G}$ with label ℓ , we assign to it a corresponding edge of $\mathcal{G}'$ from $\mathbf{w}$ to $\mathbf{w}'$ with label ℓ' given by

$$(5.5) \quad \ell' = -\ell + c \pmod{p}, \quad 0 \leq \ell' \leq p-1,$$

If the current vertex has $c = 0$ and $\ell = 0$, then the new vertex has $\ell' = 0$ and is assigned carry digit $c' = 0$. If either $c = -1$ or if $c = 0$ and $\ell > 0$, then the new carry digit $c' = -1$. (Once a path in $\mathcal{G}'$ reaches a state with carry digit $c' = -1$, all subsequent states reached have carry digit -1 .) Note that when $c = -1$ we have $-p \leq -\ell - 1 \leq -1$ and $\ell' = p - \ell - 1$.

We now let $\mathcal{G}'$ denote the part of the graph above reachable from the state $\mathbf{w}_0$. This graph has at most $2V$ vertices. We assert that $\mathcal{G}'$ is right-resolving. This is clear since the label ℓ' on exit edges from a vertex $\mathbf{w}$ are in one-one correspondence with labels on exit edges in $\mathcal{G}$ from the associated vertex v , via (5.5).

Suppose that $\mathcal{P}'$ is the path set determined by $(\mathcal{G}', \mathbf{w}_0)$, then $\mathcal{P}' = -\mathcal{P}$. We let $x_n = \sum_{j=0}^{n-1} \ell_j p^j$ and

$$y_n = \sum_{j=0}^{n-1} \ell_j p^j.$$

We have $y_n = x_n = 0$ as long as the carry digit $c = 0$. Let $\ell_r - 1$ be the first nonzero digit on the path, where the carry digit switches to -1 . From then on switches to $c = -1$, we have

$$y_n = (p - \ell_{r-1})p^{r-1} + \sum_{j=r}^{n-1} (p - \ell_j - 1)p^j = -x_n + p^n.$$

Letting $n \rightarrow \infty$ we obtain $y_\infty = -x_\infty$, establishing the result.

(4) For $k \geq 0$ the set $p^k \mathcal{P}$ consists of all sequences of $\mathcal{P}$ having k zeros adjoined at the beginning. A presentation $(\mathcal{G}', \mathbf{w})$ for this set is as follows. Let $\mathcal{G}'$ consist of $\mathcal{G}$ with the addition of k new vertices $\mathbf{w}_j$ ($0 \leq j \leq k-1$) added. Each of the new vertices has a single exit edge from $\mathbf{w}_j$ to $\mathbf{w}_{j+1}$ assigned label 0, for $0 \leq j \leq k-2$, and a similar exit edge labelled 0 from $\mathbf{w}_{k-1}$ to v_0 . The start state of $\mathcal{G}'$ is $\mathbf{w}_0$. $\square$

Remark 5.2. Theorem 5.1 excluded the case “multiplication by p^k ” with $k < 0$, which maps $\mathbb{Z}_p$ outside of $\mathbb{Z}_p$, so is not well-defined.

5.2. Proof of Theorem 1.4.

Proof of Theorem 1.4. Let r be a p -integral rational number, i.e. $\text{ord}_p(r) \geq 0$. We may factor $r = (-1)^a p^k \frac{M_1}{M_2}$, in which $a \in \{0, 1\}$, $k \geq 0$ and $\gcd(p, M_1 M_2) = 1$. Now we successively apply the constructions in Theorem 5.1 to multiply $\mathcal{P}$ by $\frac{1}{M_1}$, next multiply the resulting set by M_2 , next multiply the resulting set by $(-1)^a$, and finally multiply the resulting set by p^k . $\square$

6. EXAMPLES

In these examples, we let $\Sigma_p(D)$ denote the p -adic integer Cantor set consisting of all p -adic integers whose digits are drawn from a set $D \subseteq \{0, 1, \dots, p-1\}$. Such sets are all p -adic path sets, having Hausdorff dimension $d_H(\Sigma_p(D)) = \log_p |D|$.

Example 6.1. Consider the 3-adic Cantor set $\Sigma_3(\{0, 1\})$ whose 3-adic expansions admit the digit 2. This set is presented as a 3-adic path set by the pointed labelled graph $(\mathcal{G}, 0)$ pictured in Figure 1.




FIGURE 1. Presentation $(\mathcal{G}, 0)$ of $\Sigma_3(\{0, 1\})$.

This graph has adjacency matrix

$$\mathbf{A} = \begin{pmatrix} 2 \end{pmatrix},$$

whose Perron-Frobenius eigenvalue is 2, hence the Cantor set $\Sigma_3(\{0, 1\})$ has Hausdorff dimension $d_H(\Sigma_3(\{0, 1\})) = \log_3 2$, as is well known.

Now we consider the effect of additively shifting by $r = 2$. The construction of Section 4.1 applied to the presentation above yields the presentation of $\Sigma_3(\{0, 1\}) + 2$ given in Figure 2 by $\mathcal{G}'(0200)$.




FIGURE 2. Presentation $(\mathcal{G}', 0200)$ of $\Sigma_3^1(\{0, 1\}) + 2$.

Under one ordering of the vertices of $\mathcal{G}'$, the adjacency matrix of the underlying (undirected) graph of $\mathcal{G}'$ is

$$\mathbf{A}' = \begin{pmatrix} 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 1 \end{pmatrix}.$$

The eigenvalues of $\mathbf{A}'$ are 2 and 0 (multiplicity 3), so we see the Perron eigenvalue is 2. Thus the Hausdorff dimension is

$$d_H(\Sigma_3(\{0, 1\}) + 2) = \log_3 2.$$

Recall that these sets must have the same Hausdorff dimension, because they are bi-Lipschitz equivalent, hence their adjacency matrices must have the same Perron eigenvalue.

Example 6.2. We consider the effect of set addition on 5-adic Cantor sets $\Sigma_5(D)$ for certain subsets of digits D . For all sets of two digits, we have $d_H(\Sigma_5(\{a, b\})) = \log_5 2$. Let

$$X_{i,j} = \Sigma_5(\{0, i\}) + \Sigma_5(\{0, j\}), \text{ for } 1 \leq i, j \leq 4.$$

One can show that

$$(6.1) \quad \log_5 3 \leq d_H(X_{i,j}) \leq \log_5 4.$$

We find by inspection that the sums of certain Cantor sets are themselves Cantor sets:

$$X_{1,1} := \Sigma_5(\{0, 1\}) + \Sigma_5(\{0, 1\}) = \Sigma_5(\{0, 1, 2\})$$

$$X_{1,2} = \Sigma_5(\{0, 1\}) + \Sigma_5(\{0, 2\}) = \Sigma_5(\{0, 1, 2, 3\})$$

These examples have Hausdorff dimensions $d_H(X_{1,1}) = \log_5 3$ and $d_H(X_{1,2}) = \log_5 4$, respectively, and they show that the bounds in (6.1) are sharp. Much more interesting are the sets $X_{2,3}$ and $X_{1,4}$, which are not Cantor sets; this p -adic carry operations done on addition in the set sum destroys the Cantor set property. To compute their Hausdorff dimension, we first find p -adic path set presentations for them by the construction of Theorem 1.3. These presentations are not right-resolving, but we then apply the subset construction method in [2, Sec. 2] to obtain a right-resolving presentation. We omit the details, noting only that for $X_{1,4}$ we find the resulting graph has five vertices and adjacency matrix

$$\mathbf{A}_{14} = \begin{pmatrix} 1 & 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 & 1 \\ 1 & 1 & 0 & 1 & 1 \\ 1 & 1 & 0 & 1 & 1 \end{pmatrix}.$$


Its Perron eigenvalue is $2 + \sqrt{2}$. Computing their Hausdorff dimension by the formula of Theorem 3.1, we obtain

$$d_H(X_{1,4}) = \log_5(2 + \sqrt{2}) \approx \log_5(3.41412)$$

A similar construction for $X_{2,3}$ leads to

$$d_H(X_{2,3}) = \log_5(2 + \sqrt{3}) \approx \log_5(3.73205).$$

Example 6.3. We next consider on the effect on the 3-adic Cantor set $\Sigma_3(\{0, 1\})$ of a multiplicative translation by $r = \frac{1}{4}$. The set $\frac{1}{4}\Sigma_3(\{0, 1\})$ has a presentation $(\mathcal{H}, 00)$ obtained that of $\Sigma_3(\{0, 1\})$ given by $(\mathcal{G}, 0)$, using the construction given in Section 5.1. This presentation is shown in Figure 3.

FIGURE 3. Presentation $(\mathcal{H}, 00)$ of $\frac{1}{4}\Sigma_3(\{0, 1\})$.

The adjacency matrix $\mathbf{B}$ of the underlying graph of $\mathcal{H}$ is


$$\mathbf{B} = \begin{pmatrix} 1 & 1 & 0 & 0 \\ 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 \end{pmatrix}.$$

This matrix has Perron eigenvalue 2, and it has three other smaller nonzero eigenvalues, one real and two conjugate complex. Using the formula in Theorem 3.1(3) we obtain $d_H(\frac{1}{4}\Sigma_3(\{0, 1\})) = \log_3 2$.

Example 6.4. In this final example we consider the effect of intersecting multiplicatively translated Cantor sets. We consider the case

$$X := \Sigma_3(\{0, 1\}) \cap \frac{1}{4}\Sigma_3(\{0, 1\}).$$

We obtain by the method of [2, Sec. 4] applied to the presentations above the presentation of $\mathcal{H}'(000)$ shown in Figure 4, where $\mathcal{H}'$ is the label product $\mathcal{H}' = \mathcal{G} \star \mathcal{H}$, as defined in [2, Sec. 4].

FIGURE 4. Presentation $\mathcal{H}'(000)$ of $\Sigma_3(\{0, 1\}) \cap \frac{1}{4}\Sigma_3(\{0, 1\})$.

The adjacency matrix of the underlying graph of $\mathcal{H}'$ is

$$\mathbf{B}' = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix},$$

whose Perron eigenvalue is $\frac{1+\sqrt{5}}{2}$. We conclude that

$$d_H(\Sigma_3(\{0, 1\}) \cap \frac{1}{4}\Sigma_3(\{0, 1\})) = \log_3\left(\frac{1 + \sqrt{5}}{2}\right).$$

Remark 6.5. In [3] we will study intersections of multiplicative translates of 3-adic Cantor sets in much more detail.

REFERENCES

- [1] A. G. Abercrombie, *The Hausdorff dimension of some exceptional sets of p -adic integer matrices*, J. Number Theory **53** (1995), 311–341.
- [2] W. Abram and J. C. Lagarias, *Path sets and their symbolic dynamics*, eprint [arXiv:1207.5004](#).
- [3] W. Abram and J. C. Lagarias, *Intersections of multiplicative translates of 3-adic Cantor sets*, in preparation.
- [4] R.L. Adler and B. Marcus, *Topological entropy and equivalence of dynamical systems*, Memoirs of the American Mathematical Society, Volume 20, No. 219, AMS: Providence, RI 1979.
- [5] J.-C. Ban, W.-G. Hu and S.-S. Lin, *Pattern generation problems arising in multiplicative integer systems*, eprint [arXiv:1207.7154](#)
- [6] M. Das, Sze-Man Ngai, *Graph-directed iterated function systems with overlaps*, Indiana Univ. Math. J. **53** (2004), no. 1, 109–134.
- [7] G. Edgar, *Measure, topology and fractal geometry*, Second Edition Springer-Verlag: New York 2008.
- [8] G. E. Edgar and J. Golds, *A fractal dimension estimate for a graph-directed IFS of non-similarities*, Indiana Univ. Math. J. **48** (1999), no. 2, 429–447.
- [9] P. Erdős, *Some unconventional problems in number theory*, Math. Mag. **52** (1979), 67–70.
- [10] K. J. Falconer, *The Geometry of Fractal Sets*, Cambridge Tracts in Mathematics, No. 85, Cambridge Univ. Press: Cambridge 1985.
- [11] H. Furstenberg, *Disjointness in ergodic theory, minimal sets, and a problem in Diophantine approximation*, Math. Systems Theory **1** (1967), 1–49.
- [12] J. E. Hutchinson, *Fractals and self similarity*, Indiana Univ. Math. Journal **30** (1981), 713–747.
- [13] A. Katok and B. Hasselblatt, *Introduction to the Modern Theory of Dynamical Systems* (Cambridge University Press, New York, 1995).
- [14] R. Kenyon, Y. Peres and B. Solomyak, *Hausdorff dimension for fractals invariant under the multiplicative integers*, Ergod. Th. Dyn. Sys. **32** (2012), No. 5, 1567–1584.
- [15] J.C. Lagarias, *Ternary expansions of powers of 2*, J. London Math. Soc.(2) **79** (2009), 562–588.
- [16] D. Lind, *The entropies of topological Markov shifts and a related class of algebraic integers*, Ergod. Th. Dyn. Sys. **4** (1984), no. 2, 283–300.
- [17] D. Lind and B. Marcus, *An Introduction to Symbolic Dynamics and Coding*, (Cambridge University Press, New York, 1995).
- [18] K. Mahler, *Lectures on diophantine approximations, Part I. g -adic numbers and Roth’s theorem*, Prepared from notes of R. P. Bambah, University of Notre Dame Press, Notre Dame IN 1961.
- [19] R. D. Mauldin and M. Urbański, *Graph directed Markov systems. Geometry and dynamics of limit sets*, Cambridge Tracts in Mathematics No. 148, Cambridge Univ. Press: Cambridge 2003.
- [20] R. D. Mauldin and S. C. Williams, *On the Hausdorff dimension of some graphs*, Trans. Amer. Math. Soc. **298** (1986), no. 2, 793–803.
- [21] R.D. Mauldin and S.C. Williams, *Hausdorff Dimension of Graph Directed Constructions*, Transactions of the American Mathematical Society, **309**, No. 2 (1988) , 811–829.
- [22] S.-M. Ngai, Fei Wang and Xinhan Dong, *Graph-directed iterated function systems satisfying the generalized finite type condition*, Nonlinearity **23** (2010), 2333–2350.
- [23] Sze-Man Ngai and Y. Wang, *Hausdorff dimension of self-similar sets with overlaps*, J. London Math. Soc. **63** (2001), no. 3, 655–672.
- [24] Y. Peres, J. Schmeling, S. Seuret, B. Solomyak, *Dimensions of some fractals defined via semigroup generated by 2 and 3*, eprint: [arXiv:1206.4742](#).